

# Electrostatic self-force in a static weak gravitational field with cylindrical symmetry

B. Boisseau\*, C. Charmousis<sup>†</sup> and B. Linet<sup>‡</sup>

Laboratoire de Modèles de Physique Mathématique

CNRS/EP 93, Université François Rabelais

Faculté des Sciences et Techniques

Parc de Grandmont 37200 TOURS, France

## Abstract

We determine the electrostatic self-force for a point charge at rest in an arbitrary static metric with cylindrical symmetry in the linear approximation in the Newtonian constant. In linearised Einstein theory, we express it in terms of the components of the energy-momentum tensor.

---

\*E-mail : boisseau@celfi.phys.univ-tours.fr

<sup>†</sup>E-mail : christos@celfi.phys.univ-tours.fr

<sup>‡</sup>E-mail : linet@celfi.phys.univ-tours.fr

# 1 Introduction

DeWitt and DeWitt [1] have investigated the DeWitt-Brehme equation of motion [2] for a point charge in a static weak gravitational field with spherical symmetry. They determine the electromagnetic self-force for a point charge falling non relativistically. It is not connected with the emission of electromagnetic radiation and it is non-zero for a point charge at rest. However, for a point charge at rest in a static gravitational field within the Newtonian approximation, another approach due to Vilenkin [3] gives directly the induced electrostatic self-force. An exact determination of this in the Schwarzschild spacetime has been performed [4, 5, 6] by using the expression in closed form of the electrostatic potential [7].

A renewed interest about the electrostatic self-force has appeared in the studies of the straight cosmic strings in general relativity [8]. For an infinitely thin straight cosmic string, the determination of the electrostatic self-force has been performed by Linet [9, 10] and Smith [11]. However, some questions occur about the boundary conditions of the electrostatic field at the position of the singular source of the spacetime [12]. Therefore, electrostatics in the case of an extended straight cosmic string have been analysed [13, 14] in order to derive an expression of the electrostatic self-force.

The purpose of the present paper is to determine the electrostatic self-force for a point charge at rest in an arbitrary static metric with cylindrical symmetry. This class of spacetimes can describe extended cosmic strings in general relativity but also different types of cosmic strings eventually in other gravitational metric theories. Here, we limit ourselves to the calculation of the electrostatic self-force in the linear approximation in the Newtonian constant  $G$  (we set  $c = 1$ ). It can be derived from a self-interaction potential

energy  $W$  ; in Minkowskian-like coordinates we have

$$f^A = -\frac{\partial W}{\partial x^A} \quad A = 1, 2 \quad \text{and} \quad f^3 = 0 \quad (1)$$

We emphasize that we find the electrostatic potential beyond the Newtonian approximation in which the components of the stress tensor are neglected before the energy density. We will see that the knowledge of the electrostatic potential in the case of a straight cosmic string plays a central role in our method enabling us to obtain the result in integral form.

The plan of the paper is as follows. In Section 2, we indicate the coordinate system in which we analyse the static metrics with cylindrical symmetry. We show in Section 3 how the determination of the electrostatic potential in the linear approximation in  $G$  can be reduced in solving two equations. In Sections 4 and 5, we solve successively these equations obtaining as a consequence the self-interaction potential energy. We add some concluding remarks in Section 6.

## 2 Preliminaries

A static metric with cylindrical symmetry can be written in the coordinate system  $(x^\mu)$   $\mu = 0, 1, 2, 3$  as

$$ds^2 = g_{00}dt^2 - g_{11}[(dx^1)^2 + (dx^2)^2] - g_{33}(dx^3)^2 \quad (2)$$

where the  $g_{00}$ ,  $g_{11}$  and  $g_{33}$  depend only on  $\rho$  with  $\rho = \sqrt{(x^1)^2 + (x^2)^2}$ .

Since we seek electromagnetic effects depending on  $G$  in the first approximation, we develop the components  $g_{\mu\nu}$  of the metric (2) with respect to the Minkowskian metric  $\eta_{\mu\nu}$  as

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (3)$$

where  $h_{\mu\nu}$  are the linear terms in  $G$ . According to (2), we see that

$$h_{11} = h_{22} \quad \text{and} \quad h_{0i} = 0 \quad (4)$$

The linearised Einstein equations for  $h_{\mu\nu}$  take the following form in which we must take into account the staticity and the cylindrical symmetry

$$\Delta h_{\mu\nu} + \partial_\mu(\eta^{\sigma\rho}\partial_\sigma h_{\nu\rho} - \frac{1}{2}\partial_\nu h) + \partial_\nu(\eta^{\sigma\rho}\partial_\sigma h_{\mu\rho} - \frac{1}{2}\partial_\mu h) = 16\pi G(T_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}T) \quad (5)$$

where  $h = \eta^{\mu\nu}h_{\mu\nu}$ ,  $\Delta$  being the Laplacian operator which reduces here to  $\Delta_2$  the Laplacian operator in two dimensions. The energy-momentum tensor  $T_{\mu\nu}$  which is the source of the gravitational field is the one calculated for a material distribution in the Minkowskian spacetime. It is assumed to be regular.

The metric of an extended cosmic string is a particular case of metric (2). In the limit case where its radius tends to zero, we get the conical metric with a singular source. In the linear approximation in  $G$ , the conical metric can be written in the present coordinates [15] as

$$ds^2 = (dx^0)^2 - (1 - 8G\mu \ln \frac{\rho}{\rho_*})[(dx^1)^2 + (dx^2)^2] - (dx^3)^2 \quad (6)$$

where  $\mu$  is the linear mass density of the cosmic string. We have introduced in (6) an arbitrary length  $\rho_*$  which can eventually represent the radius of the cosmic string.

### 3 Equation of the electrostatic potential

For electrostatics in a static metric, we use the electrostatic potential  $V$  to which the vector potential  $A_\mu$  reduces in this case

$$A_0 = V \quad \text{and} \quad A_i = 0 \quad (7)$$

and it does not depend on the coordinate  $x^0$ . From the Maxwell equations in the background metric (2), we obtain the equation of the electrostatic potential

$$\partial_i[\sqrt{-g}g^{ij}g^{00}\partial_j V] = \frac{\sqrt{-g}}{\epsilon_0}j^0 \quad (8)$$

where  $j^0$  is the charge density. For a point charge  $q$  located at  $x^i = x_0^i$ , we have

$$j^0(x^i) = \frac{q}{\sqrt{-g}}\delta^{(3)}(x^i - x_0^i) \quad (9)$$

The electrostatic potential  $V$  has to be symmetric in the variables  $x^i$  and  $x_0^i$ .

We can introduce development (3) of the metric (2) in equation (8). It is convenient to use the following notations

$$g^{00} = 1 + h^{00} \quad \text{and} \quad \sqrt{-g}g^{ij} = -\delta^{ij} + \gamma^{ij} \quad (10)$$

where  $h^{00}$  and  $\gamma^{ij}$  are linear terms in  $G$  given by

$$\gamma^{ij} = h^{ij} - \frac{1}{2}h\delta^{ij} \quad \text{and} \quad h^{\mu\nu} = -\eta^{\mu\sigma}\eta^{\nu\rho}h_{\sigma\rho} \quad (11)$$

It will be useful for determining the electrostatic potential to recast  $\gamma^{ij}$  in the form

$$\gamma^{ij} = \gamma_C^{ij} + \gamma^{11}\delta^{ij} \quad \text{with} \quad \gamma_C^{ij} = \delta_3^i\delta_3^j(\gamma^{33} - \gamma^{11}) \quad (12)$$

which is possible taking into account (4). We remark that metric (6) corresponds to  $h^{00} = 0$  and  $\gamma_C^{ij} = \gamma_{CS}^{ij}$  where  $\gamma_{CS}^{ij}$  is defined by  $\gamma_{CS}^{33} = 8G\mu \ln \rho/\rho_*$ .

According to (12), equation (8) with source term (9) can be now written in the following manner

$$\partial_i[(-\delta^{ij} - \delta^{ij}(h^{00} - \gamma^{11}) + \gamma_C^{ij})\partial_j V] = \frac{q}{\epsilon_0}\delta^{(3)}(x^i - x_0^i) \quad (13)$$

In order to solve (13), we write down  $V$  as the sum of the Coulombian potential and terms linear in  $G$ . We set

$$V(x^i, x_0^i) = \frac{q}{4\pi\epsilon_0 r(x^i, x_0^i)} + V_U(x^i, x_0^i) + V_C(x^i, x_0^i) \quad (14)$$

where  $r(x^i, x_0^i)$  is the Euclidean distance between the points  $x^i$  and  $x_0^i$ . If  $V_U$  and  $V_C$  obey respectively the following equations

$$\Delta V_U + \frac{q}{4\pi\epsilon_0} \partial_i [(h^{00} - \gamma^{11}) \delta^{ij} \partial_j \frac{1}{r}] = 0 \quad (15)$$

$$\Delta V_C - \frac{q}{4\pi\epsilon_0} \partial_i [\gamma_C^{ij} \partial_j \frac{1}{r}] = 0 \quad (16)$$

then equation (13) is satisfied. We recall that we must find  $V_U$  and  $V_C$  symmetric in the variables  $x^i$  and  $x_0^i$ .

For a straight cosmic string, the exact electrostatic potential is known in another coordinate system in which the metric is conical. A local expression is given for  $G\mu < 1/8$  [10] as the sum of the Coulombian potential and a regular part assuming that point  $x$  is near point  $x_0$ . At the linear approximation in  $G$ , a change of coordinate system to obtain form (6) of the metric yields simply

$$V(x^i, x_0^i) = \frac{q}{4\pi\epsilon_0 r(x^i, x_0^i)} + V^*(x^i, x_0^i) \quad (17)$$

valid only for a point  $x^i$  near the point  $x_0^i$  where the part  $V^*$  is regular at  $x^i = x_0^i$ . It furnishes a solution  $V_C = V^*$  to equation (16) for  $\gamma_C^{ij} = \gamma_{CS}^{ij}$ . The self-interaction potential energy  $W_{CS}$  is then given by

$$W_{CS}(x_0^A) = \frac{q}{2} V^*(x_0^i, x_0^i) \quad (18)$$

which does not depend on  $x_0^3$ . It has been explicitly calculated and it does not depend on the coordinate system at the linear approximation in  $G$ . We

have

$$W_{\text{CS}}(x_0^A) = \frac{q^2 G \mu}{16 \epsilon_0 \rho_0} \quad (19)$$

As expected, it is a function of  $\rho_0$  with  $\rho_0 = \sqrt{(x_0^1)^2 + (x_0^2)^2}$ .

## 4 Electrostatic self-force induced by $V_{\text{U}}$

We follow the method of Vilenkin [3] for determining  $V_{\text{U}}$ . We seek the solution to equation (15) in the form

$$\begin{aligned} V_{\text{U}}(x^i, x_0^i) = & -\frac{q}{8\pi\epsilon_0} \left( h^{00}(x_0^i) - \gamma^{11}(x_0^i) \right) \frac{1}{r(x^i, x_0^i)} \\ & -\frac{q}{8\pi\epsilon_0} \left( h^{00}(x^i) - \gamma^{11}(x^i) \right) \frac{1}{r(x^i, x_0^i)} \\ & + \psi(x^i, x_0^i) \end{aligned} \quad (20)$$

Then, the quantity  $\psi$  satisfies

$$\Delta \psi = \frac{q}{8\pi\epsilon_0 r(x^i, x_0^i)} \Delta(h^{00} - \gamma^{11}) \quad (21)$$

The solution to equation (21) is

$$\psi(x^i, x_0^i) = -\frac{q}{32\pi^2\epsilon_0} \int \frac{\Delta(h^{00} - \gamma^{11})(b^i)}{r(b^i, x_0^i)r(b^i, x^i)} d^3b \quad (22)$$

which is symmetric in  $x^i$  and  $x_0^i$ .

From (22), we immediately obtain the self-interaction potential energy

$$W_{\text{U}}(x_0^i) = \frac{q}{2} \psi(x_0^i, x_0^i) \quad (23)$$

which is regular in  $x^i = x_0^i$ . It is given by the integral

$$W_{\text{U}}(x_0^i) = -\frac{q^2}{64\pi^2\epsilon_0} \int \frac{\Delta(h^{00} - \gamma^{11})(b^i)}{r^2(b^i, x_0^i)} d^3b \quad (24)$$

which is well defined.

In the case of the cylindrical symmetry,  $\Delta$  acting on  $h^{00} - \gamma^{11}$  is in fact  $\Delta_2$  the Laplacian operator in two dimensions. Moreover, we can perform in (24) the integration on the variable  $b^3$  and we obtain

$$W_U(x_0^A) = -\frac{q^2}{64\pi\epsilon_0} \int \frac{\Delta_2(h^{00} - \gamma^{11})(b^A)}{\rho(x_0^A, b^A)} d^2b \quad (25)$$

where  $\rho(x_0^A, b^A) = \sqrt{(x_0^1 - b^1)^2 + (x_0^2 - b^2)^2}$ . As a result of the integration on  $b^3$ ,  $W_U$  does not depend on  $x_0^3$ . Furthermore, it is easy to see that  $W_U$  is a function of  $\rho_0$ .

In general relativity, we may use the linearised Einstein equations (5) having as source term the energy-momentum tensor  $T_{\mu\nu}$  to calculate  $\Delta_2(h^{00} - \gamma^{11})$ . After some manipulations, we find

$$W_U(x_0^A) = \frac{q^2 G}{8\epsilon_0} \int \frac{(T_0^0 - T_3^3)(b^A)}{\rho(x_0^A, b^A)} d^2b \quad (26)$$

## 5 Electrostatic self-force induced by $V_C$

We now turn our attention to determining  $V_C$ . We can solve equation (16) if we know the solution  $G(x^i, x_0^i; b^A)$  to the following equation

$$\Delta G - \frac{q}{4\pi\epsilon_0} \partial_i \left[ \delta_3^i \delta_3^j \delta^{(2)}(x^A - b^A) \partial_j \frac{1}{r(x^i, x_0^i)} \right] = 0 \quad (27)$$

the desired solution will be written in an integral form

$$V_C(x^i, x_0^i) = \int \gamma_C^{33}(b^A) G(x^i, x_0^i; b^A) d^2b \quad (28)$$

We have already noticed that  $V^*$ , appearing in (17) for a straight cosmic string, satisfies (16) which in this particular case takes the form

$$\Delta V^* - \frac{2G\mu q}{\pi\epsilon_0} \partial_i \left[ \delta_3^i \delta_3^j \ln \frac{\rho}{\rho_*} \partial_j \frac{1}{r(x^i, x_0^i)} \right] = 0 \quad (29)$$



As mentioned this solution is known for  $x^i$  near  $x_0^i$ . A change of origin of the coordinates ( $x^A$ ) by  $b^A$  in equation (29) yields the equation

$$\Delta V_b^* - \frac{2G\mu q}{\pi\epsilon_0} \partial_i \left[ \delta_3^i \delta_3^j \ln \frac{\rho(x^A, b^A)}{\rho_*} \partial_j \frac{1}{r(x^i, x_0^i)} \right] = 0 \quad (30)$$

whose solution  $V_b^*$  is consequently given by

$$V_b^*(x^i, x_0^i; b^A) = V^*(x^A - b^A, x^3; x_0^A - b^A, x_0^3) \quad (31)$$

But  $\Delta_2 \ln \rho / \rho_* = 2\pi\delta^{(2)}$ , therefore

$$G(x^i, x_0^i; b^A) = \frac{1}{16\pi G\mu} \Delta_2 V_b^*(x^i, x_0^i; b^A) \quad (32)$$

where  $\Delta_2$  acts on the variables  $b^A$ . According to (28), we are now in a position to give the integral expression of the desired solution

$$V_C(x^i, x_0^i) = \frac{1}{16\pi G\mu} \int \Delta_2 \gamma_C^{33}(b^A) V_b^*(x^i, x_0^i; b^A) d^2b \quad (33)$$

for a point  $x^i$  near point  $x_0^i$ , in which we have used a property of the convolution product so that  $\Delta_2$  applies now on  $\gamma_C^{33}$ . Expression (33) is regular at  $x^i = x_0^i$ .

From (33), we deduce immediately the self-interaction potential energy

$$W_C(x_0^A) = \frac{q}{32\pi G\mu} \int \Delta_2 \gamma_C^{33}(b^A) V_b^*(x_0^i, x_0^i; b^A) d^2b \quad (34)$$

By considering the self-interaction potential energy derived from (19) for a straight cosmic string located at  $x^A = b^A$

$$\frac{1}{2} q V_b^*(x_0^i, x_0^i; b^A) = \frac{q^2 G\mu}{16\epsilon_0 \rho(x_0^A, b^A)} \quad (35)$$

we get from (34) an integral expression

$$W_C(x_0^A) = \frac{q^2}{256\pi\epsilon_0} \int \frac{\Delta_2 \gamma_C^{33}(b^A)}{\rho(x_0^A, b^A)} d^2b \quad (36)$$

In general relativity we may use the linearised Einstein equations (5) to calculate  $\Delta_2 \gamma_C^{33}$ ; we obtain after some manipulations

$$W_C(x_0^A) = \frac{q^2 G}{16\epsilon_0} \int \frac{(T_3^3 - T_1^1 - T_2^2)(b^A)}{\rho(x_0^A, b^A)} d^2 b \quad (37)$$

## 6 Conclusion

For a static spacetime with cylindrical symmetry, we have determined in the linear approximation in  $G$  the self-interaction potential energy  $W_U + W_C$  given respectively by (25) and (36) in terms of the components of the metric. We can apply these formulas for all gravitational metric theories.

Taking into account the linearised Einstein equations, we have expressed these potentials with the aid of the components of the energy-momentum tensor which is the source of the gravitational field. We have found the following self-interaction potential energy

$$W(x_0^A) = \frac{q^2 G}{8\epsilon_0} \left[ \int \frac{T_0^0(b^A)}{\rho(x_0^A, b^A)} d^2 b - \frac{1}{2} \int \frac{(T_3^3 + T_1^1 + T_2^2)(b^A)}{\rho(x_0^A, b^A)} d^2 b \right] \quad (38)$$

which depends only on  $\rho_0$ . For an energy-momentum tensor vanishing at infinity, we point out that the integrals in expression (38) are well defined. We emphasize that formula (38) can be now evaluated in arbitrary Minkowskian-like coordinates in the linear approximation in  $G$ .

Formula (38) is general for regular energy-momentum tensors. It can be applied for different types of cosmic string. This formula obviously verifies the limit case in which the radius of an extended cosmic string tends to zero since

$$T_0^0 \rightarrow \mu \delta^{(2)} \quad T_3^3 \rightarrow \mu \delta^{(2)} \quad T_1^1 \rightarrow 0 \quad T_2^2 \rightarrow 0 \quad (39)$$

giving expression (19) for an infinitely thin cosmic string.

## References

- [1] DeWitt C M and DeWitt B S 1964 *Physics* **1** 3
- [2] DeWitt B S and Brehme R W 1960 *Ann. Phys. (N.Y.)* **9** 220
- [3] Vilenkin A 1979 *Phys. Rev D* **20** 373
- [4] Smith A G and Will C M 1980 *Phys. Rev. D* **22** 1276
- [5] Zel'nikov A J and Frolov V P 1982 *Sov. Phys. JETP* **55** 191
- [6] Léauté B and Linet B 1983 *Int. J. Theor. Phys.* **22** 67
- [7] Linet B 1976 *J. Phys. A : Math. Gen.* **9** 1081
- [8] Vilenkin A 1986 “*Three Hundred Years of Gravitation*” ed S W Hawking and W Israel (Cambridge : Cambridge University Press) p 499
- [9] Linet B 1986 *Phys. Rev. D* **33** 1833
- [10] Linet B 1986 *Ann. Inst. H. Poincaré - Phys. Théor.* **45** 249
- [11] Smith A G 1990 “*The Formation and Evolution of Cosmic Strings*” ed G W Gibbons, S W Hawking and T Vaschupati (Cambridge : Cambridge University Press) p 262
- [12] Kay B S and Studer U M 1991 *Commun. Math. Phys.* **139** 103
- [13] Perkins W B and Davis A-C 1991 *Nucl. Phys. B* **349** 207
- [14] Allen B and Ottewill A C 1990 *Phys. Rev. D* **42** 2669
- [15] Vilenkin D 1981 *Phys. Rev. D* **23** 852